arXiv:math/0407087v1 [math.AG] 6 Jul 2004

ON THE 0-DIMENSIONAL IRREDUCIBLE COMPONENTS OF THE SINGULAR LOCUS OF $\mathcal{A}_g$

V. González -Aguilera, J.M. Muñoz-Porras and Alexis G. Zamora

Abstract. In this short note we give a characterization of extremal principally polarized abelian varieties determining an isolated point in $Sing\mathcal{A}_g$. The case $g=5$ is treated with detail.

1. Introduction

Let $\mathcal{A}_g$ be the moduli space of principally polarized abelian varieties (p.p.a.v.). In [3] it was proved the existence of irreducible varieties $\mathcal{A}_g(p,\rho_n)\subset\mathcal{A}_g$ representing triples (X,σ,α) consisting of a p.p.a.v. X, a non trivial automorphism $\sigma\in Aut(X)$ of order p (a prime number) and the conjugation class α of the representation of σ in X_n for some fixed $n\geq 3$. The local information about these varieties is encoded in the so called "analytic representation" of σ, that is, the differential

$$d\sigma : T_0X\longrightarrow T_0X.$$

Indeed, if $\{\xi^{i_1},...,\xi^{i_g}\}$ denotes the set of eigenvalues of $d\sigma$ then

$$\dim\mathcal{A}_g(p,\rho_n)=\frac{n_0(n_0+1)}{2}+\sum_{i=1}^{p-1/2}n_in_{p-i},$$

where n_i denotes the dimension of the eigenspace associated to ξ^i. Moreover, there exists a number r such that $n_i+n_{p-i}=r$ for $i\neq 0$ and

$$g=n_0+\frac{p-1}{2}r.$$

The existence of such number is a consequence of the fact that the rational representation of σ is the direct sum $d\sigma\oplus\bar{d\sigma}$ ([3], Prop.2.1).

Two fundamental problems arise from these results. In the first place there is a problem of existence: given numerical data satisfying the above restriction, it there exists an p.p.a.v. X admitting an automorphism compatible with this data? In the second place, in order to describe the irreducible components of $Sing\mathcal{A}_g$ we need to determine what inclusions of the form $\mathcal{A}_g(p,\rho_n)\subset\mathcal{A}_g(q,\rho'_n)$ occurs. As we shall show both problems are closely related.

1991 *Mathematics Subject Classification*. 14K10, 14K22.

The first author was partially supported by Fondecyt Grant N1010432 and DGIP UTFSM Grant 120321. The second author was partially supported by MCT Grant BFM 2003-00078 and JCyL Grant SA071/04. The third author was partially supported by Conacyt Grant 41459-F.

Typeset by $\mathcal{AMS}$-TEX

The first was solved positively in [3] in the case $n_0 = 0$, the idea is to reduce the existence problem to the existence of p.p.a.v. of dimension g admitting an automorphism of prime order $= 2g+1$ (we shall call such a p.p.a.v. an extremal one), note that in this case $r = 1$, $n_0 = 0$ and $n_i n_{p-i} = 0$. The existence of these varieties is ensured by a classical construction ([5], section 22, [L], Chapter 1) which in fact is valid in a more general context: the existence of CM-type Abelian Varieties. Then the existence problem is solved by considering products of these "extremal varieties" ([3], Lemma 2.7).

In this paper we study and solve the problem of inclusion of varieties $\mathcal{A}_g(p, \rho_n) \subset \mathcal{A}_g(q, \rho'_n)$ in the extremal case, i.e. under the assumption $p = 2g+1$. It should be noticed that if the number $2g+1$ is a prime, then it is the maximal possible order for an automorphism of prime order acting on p.p.a.v. of dimension g. It is clear that $\dim \mathcal{A}_g(2g+1, \rho) = 0$.

In order to fix the notation we recall the construction of extremal p.p.a.v. The starting point is $\mathbb{F}_p^*$ which we identify with $\{\xi, \xi^2, ..., \xi^{p-1}\}$, next, fix a partition $\{C, C'\}$ of $\mathbb{F}_p^*$ with $|C| = g$. Let $C = \{\xi^{i_1}, ..., \xi^{i_g}\}$. Define the isomorphism:

$$\Phi_C : \mathbb{Q}(\xi) \otimes_{\mathbb{Q}} \mathbb{R} \longrightarrow \mathbb{C}^g,$$

given by $\Phi_C(\xi) = (\xi^{i_1}, ..., \xi^{i_g})$. Obviously ξ acts on $\mathbb{C}^g$ as a linear automorphism of order p under this identification. Let I be a fractional ideal of $\mathbb{Q}(\xi)$. $L := \Phi_C(I) \subset \mathbb{C}^g$ is a lattice invariant under the action of ξ. In this way $X = \mathbb{C}^g/L$ is a complex torus admitting an action of $\mathbb{Z}_p$ with analytic representation determined by the set of eigenvalues C. A Riemann form can be defined on $(\mathbb{C}^g, L)$ in such a way that X becomes an extremal p.p.a.v.

Multiplication defines a natural action of $\mathbb{F}_p^*$ in the set $\mathcal{C}$ of all the subsets of $\mathbb{F}_p^*$ of cardinality g. Let H_C be the isotropy group of C under this action. The main result of this note is the following:

Theorem 1. . *Let X be an extremal p.p.a.v.. Denote by σ the automorphism of X of order $2g+1$. Assume that the analytic representation $d\sigma$ has as set of eigenvalues the set C. Then X is an isolated point in $Sing\mathcal{A}_g$ if and only if H_C is trivial.*

Section 2 of the paper is devoted to prove this theorem. After this we include in section 3 a classification of the extremal p.p.a.v. of dimension 5 that determine isolated points in $SingA_5$. The proof of Theorem 1 is based in our previous paper [3] as well as in the results in [2].

2. Proof of Theorem 1

Proof of Theorem 1.

Following the notation introduced in the previous section, let X be an extremal p.p.a.v. constructed from the set C. $\mathbb{F}_p^*$ is the Galois group of $\mathbb{Q}(\xi)/\mathbb{Q}$, an element k of $\mathbb{F}_p^*$ acts in $\mathbb{Q}(\xi)$ by means of $k.\xi = \xi^k$. The relevance of the group of isotropy $H_C \subset \mathbb{F}_p^*$ defined in the introduction lies in the following observation: let $\theta \in H_C$, $\theta \neq 1$. θ can be extended to a $\mathbb{R}$-linear automorphism:

$$\theta : \mathbb{R}(\xi) \longrightarrow \mathbb{R}(\xi),$$

as $\theta C = C$ the map $\Phi_C(\theta)$ is a $\mathbb{C}$-linear isomorphism. Furthermore, θ determines an element of $Aut(X)$, that we are going to denote for simplicity just by θ. In this way $H_C \neq 1$ implies that the total group of automorphism of X contains two different cyclic subgroups: $< \xi >$ and $< \theta >$. Call G to the total automorphism group, $G := Aut(X)$, and $G_+ := G/\{\pm 1\}$.

We have the following exact sequence of groups:

$$1 \to Z_G(< \xi >) \to N_G(< \xi >) \xrightarrow{\phi} H_C \to 1, \tag{1.1}$$

with $N_G(< \xi >)$ (respectively $Z_G(< \xi >)$ denoting as usual the normalizer (resp. the centralizer) of $< \xi >$ in G. The only map that needs definition is ϕ, if $g \in N_G(< \xi >)$, $g.\xi.g^{-1} = \xi^k$, then we define $\phi(g) = k \in \mathbb{F}_p^*$. ϕ is surjective, as ξ and ξ^k are conjugate if and only if they have the same set of eigenvalues. Note that, by construction, $\theta\xi\theta^{-1} = \xi^\theta$.

The sequence 1.1 induces the following:

$$1 \to < \xi > \to N_+ \xrightarrow{\phi_+} H_C \to 1, \tag{1.2}$$

where $N_+ = N_G(< \xi >)/\pm 1$. In fact, $\phi_+(g) = 1$ means $g\xi = \xi g$. Now, g as an automorphism of $\mathbb{Q}(\xi)$ must acts as multiplication for some element: $g(x) = ux$, $\forall x \in \mathbb{Q}(\xi)$, and, as g commutes with ξ, $uI = I$. It follows that $u \in \mathbb{Q}(\xi)^*$ and $g \in < \xi >$.

Now, we can conclude the proof of the theorem: if $H_C = 1$ then by the previous exact sequence $N_+ = < \xi >$ and by Theorem 2 in [2] $G_+ = < \xi >$.

Conversely, assume $H_C \neq 1$ and let $\theta \in H_C$ be an element of order q, a divisor of $(p-1)$. The action of θ on $\mathbb{Q}(\xi)$ gives rise to a decomposition:

$$\mathbb{Q}(\xi) \simeq \mathbb{Q}(\xi)^{H_C} \oplus M,$$

where $\mathbb{Q}(\xi)^{H_C}$ is the module of H_C- invariants and M its complement. Moreover, $\mathbb{Q}(\xi)^{H_C} \neq 0$, since $\theta(\xi^{k_1} + ... \xi^{k_g}) = \xi^{k_1} + ... + \xi^{k_g}$. After tensoring with $\mathbb{R}$ we obtain:

$$V \simeq V_0 \oplus ... \oplus V_m,$$

with $V_0 = \mathbb{Q}(\xi)^{H_C} \otimes \mathbb{R} \neq 0$. We conclude that the space of θ-invariants V_0 is of strictly positive dimension. Thus, $X \in \mathcal{A}_g(q, \theta_n)$ with

$$\dim \mathcal{A}_g(q, \theta_n) \geq 1.$$

□

The previous characterization gives, moreover, a nice classification of extremal p.p.a.v. in terms of the simplicity of these varieties. This is provided by a theorem of Shimura and Taniyama ([6], [4]). Explicitly:

Corollary 1.1. *Let X be an extremal p.p.a.v.. X is an isolated point of $Sing\mathcal{A}_g$ if and only if X is a simple Abelian variety.*

Proof. Just combine Theorem 1 with Theorem 3.5, Chapter 1 in [4]. □

Another useful remark is that there exists a simple sufficient condition that guarantees the existence of extremal p.p.a.v. with $H_C = 1$. In fact, if $C = \{k_1, ..., k_g\}$ and $\theta C = C$ then

$$k_1 + ... + k_g \cong \theta(k_1 + ... + k_g) \mod (p),$$

and, therefore, $k_1 + ... + k_g$ is divisible by p. In particular, for each $g = (p-1)/2$, extremal Abelian varieties constructed from $C = \{1, ..., g\}$ are always isolated points in $Sing\mathcal{A}_g$. The argument is taken from [6], 8.4, page 72.

3. Extremal Abelian Varieties of dimension 5

In this sections we work out with some detail the case $g = 5$, $p = 11$. First of all, the number of different (non-equivalent) partitions C of $\mathbb{F}_{11}^*$ is 4, they can be represented by:

$$C_1 = \{\xi, \xi^2, \xi^3, \xi^4, \xi^5\}, \quad C_2 = \{\xi, \xi^2, \xi^3, \xi^4, \xi^6\},$$

$$C_3 = \{\xi, \xi^2, \xi^3, \xi^5, \xi^7\}, \quad C_4 = \{\xi, \xi^3, \xi^4, \xi^5, \xi^9\}.$$

Denote by X_i the extremal p.p.a.v. corresponding to each of these partitions. These variety are uniquely determined by C_i, since $\mathbb{Z}[\xi]$ is a principal ideal domain if ξ is a root of 11. We have:

Theorem 2. *The only extremal p.p.a.v. in $\mathcal{A}_5$ which are isolated in $Sing\mathcal{A}_5$ are X_1, X_2 and X_3. Moreover:*

i) X_1 is isomorphic to the Jacobian variety of the hyperelliptic curve $y^2 = \prod_{j=1}^{11}(x - \xi^j)$.

ii) X_2 is isomorphic to the Jacobian variety of the Lefschetz curve $y^{11} = x^2(x-1)^8$.

iii) X_4 (which is not isolated) is isomorphic to the Jacobian variety of the Klein's smooth hypersurface of degree 3 in $\mathbb{P}^4$.

Proof. The first assertion follows from an explicit computation: only C_1, C_2 and C_3 have trivial isotropy group and we can apply Theorem 1.

Assertions i) and ii) follows from the computation of explicit basis of $H^0(Y, \omega_Y)$ (Y denoting the corresponding curve). In these spaces $\mathbb{Z}_{11}$ acts in a natural way, it is a simple calculus to show that the analytic spectrum of these action correspond, respectively to C_1 and C_2. For instance, for i), a basis for $H^0(Y, \omega_Y)$ if Y is given by the above equation is

$$\{x^{i-1}\frac{dx}{y} \mid i = 1, ..., g\},$$

a simple computation shows that $\mathbb{Z}_{11}$ acts on this basis by permutation, and the induced representation is given by a diagonal matrix with set of eigenvalues equal to C_1. Analogously, a basis for the space of differentials on the Lefschetz curve can be obtained from adjunction, and a similar calculus on the action of $\mathbb{Z}_{11}$ leads to the desired conclusion.

Finally, part iii) is a consequence of [1]. Note that in this case X_4 admits $PSL(2, \mathbb{F}_{11})$ as the complete group of automorphism (compare with [2], Thm. 2).

We remark that X_4 gives another example of an inclusion of the type $\mathcal{A}_g(p, \alpha) \subset \mathcal{A}_g(q, \beta)$, this time with $g = 5$, $p = 11$ and $q = 5$.

REFERENCES

1. Adler, A., *Some Integral Representations of* $PSL(2, \mathbb{F}_{11}^*)$ *and their Applications*, Jour. of Algebra **72** (1981), 115–145.
2. Bennama, H. and Bertin, J., *Remarques sur les varietiés abéliennes avec un automorphisme d'ordre premier*, Manuscripta Math. **94** (1997), 409–425.
3. González-Aguilera, V. Muñoz-Porras, J.M. and Zamora, A. G., *On the irreducible components of the singular locus of* $\mathcal{A}_g$, Jour. of Algebra **240** (2001), 230–250.
4. Lang, S., *Complex Multiplication*, Springer Verlag, 1983.
5. Mumford, D., *Abelian Varieties*, Oxford University Press, 1974.
6. Shimura, G. and Taniyama, Y., *Complex multiplication of abelian varieties and its applications to number theory*, Math. Soc. Japan, 1961.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD SANTA MARÍA,
AVENIDA ESPAÑA 1680
VALPARAÍSO, CHILE.
E-mail address: `victor.gonzalez@ usm.cl`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE SALAMANCA
PLAZA DE LA MERCED 1-4,
SALAMANCA 37008, ESPAÑA
E-mail address: `jmp@ usal.es`

CIMAT,
CALLEJÓN DE XALISCO S/N
VALENCIANA, GUANAJUATO, GTO.,
C.P. 36240,
MÉXICO
E-mail address: `alexis@ cimat.mx`